



Acer xfreemanii 'Armstrong'



Výška	10 -15 m
Šírka	4-6m
Koruna	columnar, dark, dense crown
Kôra a vetvy	no particulars
Listy	3/5 lobes, green, 10 -20 cm
Jesenné sfarbenie	Oranžová, Červená
Kvety	flower corymbs, red, March/April
Plody	none, fruitless cultivar
Tríne	Žiadny
Jedovatosť	Zvyčajne nie je jedovaté pre ľudí a väčšie zvieratá
Typ pôdy	not too limy
Spevnené plochy	Toleruje spevnenie
Zóna odolnosti	4 (-34,4 do -28,9 °C)
Odolnosť voči vetru	very good
Ostatná odolnosť	Odolné mrazu (WH 1–6), Odolné vetru
Stromy pre zvieratá a hmyz	Odolné mrazu (WH 1–6), Odolné vetru, Cenné pre motýle
Použitie	bulváre a široké ulice, úzke ulice
Tvar	Vysokokmeň, Viackmeň
Pôvod	Scanlon Nurseries, Ohio, USA, 1951
Synonymum	Acer rubrum 'Armstrong'

This cultivar originated in the USA in 1951 and until recently was considered to belong to A. rubrum. Recent studies reveal that 'Armstrong' came into being from a crossing between A. rubrum and A. saccharinum and therefore belongs to the hybrid Acer x freemanii. This hybrid was not only created by man, for it also grows in the wild in the USA. The green leaves have 3 to 5 lobes. Their under surface is grey-green. In our country the beautiful orange-red autumn colour can only be seen after a hot summer. It roots widely with a dense branching. 'Armstrong' requires a not too limy soil. The tree can be used well in hard surfaces, which renders it very suitable to be used in narrow streets and avenues.