



## Acer triflorum



<b>Výška</b>	10 m
<b>Šírka</b>	6-8m
<b>Koruna</b>	round, half-open crown
<b>Kôra a vetvy</b>	young twigs yellowish brown, older bark brownish grey, slightly scaling
<b>Listy</b>	trifoliate, green, 3 - 7 cm
<b>Jesenné sfarbenie</b>	Oranžová, Červená, Fialová
<b>Kvety</b>	yellow, in smalls corymbs of three, April/May
<b>Plody</b>	winged nutlets, 4 - 5 cm long, hairy, 4 - 5 cm
<b>Tríne</b>	Žiadny
<b>Jedovatosť</b>	Zvyčajne nie je jedovaté pre ľudí a väčšie zvieratá
<b>Typ pôdy</b>	neutral to slightly acidic
<b>Spevnené plochy</b>	Netoleruje spevnené plochy
<b>Zóna odolnosti</b>	6 (-23,3 do -17,8 °C)
<b>Odolnosť voči vetru</b>	good
<b>Ostatná odolnosť</b>	Odolné mrazu (WH 1–6), Odolné vetru
<b>Použitie</b>	stromy do nádob, strešné záhrady, malé záhrady, dvory
<b>Tvar</b>	Vysokokmeň, Viackmeň
<b>Pôvod</b>	Northern China, Korea

Bush to large tree, rarely medium-size. Somewhat resembles *A. griseum* at first sight, but differs from this in (among other things) its brown-grey, only slightly scaling bark. With *A. griseum*, the bark is reddish brown and profusely scaling. The leaves of *A. triflorum* are hairy and rather more blue-green on the underside. In summer they are green, but in autumn they turn a splendid orange to purplish red. The species name is derived from the fact that each flower corymb comprises only three flowers. *A. triflorum* should mainly be used as a solitary tree in a garden or park. It prefers a slightly acidic soil.