

https://www.vdberk.sk/index.php/stromy/betula-costata/



Betula costata



12 – 25 (30) m
10-15m
broad ovoid to round, half-open crown
light grey-brown to greyish yellow, flaking
ovoid to narrow ovoid, green, 5 - 10 cm
catkins, ? single and erect, ? 4 - 6 cm, April
ovoid fruiting catkins, 2 - 3 cm
Žiadny
Zvyčajne nie je jedovaté pre ľudí a väčšie zvieratá
few soil requirements, not drought-tolerant
Netoleruje spevnené plochy
5a (-28,8 do -26,1 °C)
moderate
Odolné mrazu (WH 1–6)
Odolné mrazu (WH 1–6), Cenné pre motýle
Vysokokmeň, Viackmeň
Eastern and central China, Korea, Manchuria

Large tree with a broad ovoid to rounded, semi-open crown. Needs a lot of space due to its broad, fairly low branching pattern. Has attractively flaking branches and bark. Not to be confused with types of B. ermanii, especially 'Blush', which is still being grown as B. costata. One difference is that the bark of B. costata already starts to peel in the second or third year, while that of B. ermanii only starts when mature. Another is that the first-year twigs of B. costata are rough and those of B. ermanii are smooth. Finally, the leaves of B. costata are more than twice as long as they are wide, and have a straight leaf base. The leaves of B. ermanii are shorter and have a somewhat heart-shaped leaf base. Has poor tolerance for drought and is sensitive to (sea) wind.