



Carya laciniosa



| | |
|-----------------------------------|---|
| Výška | 25 - 30 m |
| Šírka | 12-20m |
| Koruna | conical to rounded, open, half-open crown |
| Kôra a vetvy | trunk grey, flaking off in long pieces, twigs with felt-like hair |
| Listy | imparipinnate with 5 to 9 leaves, light green, 10 - 22 cm, feathered leaf |
| Jesenné sfarbenie | Žltá |
| Kvety | ? catkins approx. 12 - 20 cm, ? flowers in short ears, May |
| Plody | round nut, Ø approx 4 - 5 cm, husk up to 7 cm, edible, edible fruits |
| Tríne | Žiadny |
| Jedovatosť | Zvyčajne nie je jedovaté pre ľudí a väčšie zvieratá |
| Typ pôdy | humid and rich |
| Spevnené plochy | Netoleruje spevnené plochy |
| Zóna odolnosti | 6a (-23,3 do -20,6 °C) |
| Odolnosť voči vetru | good |
| Ostatná odolnosť | Odolné mrazu (WH 1–6), Odolné vetru |
| Stromy pre zvieratá a hmyz | Odolné mrazu (WH 1–6), Odolné vetru, Cenné pre včely (medonosné) |
| Tvar | Vysokokmeň |
| Pôvod | eastern part of the United States |

A tall tree with relatively thin, strongly spreading branches and twigs. Grows up to 50 m in height in its natural habitat but is practically always smaller when cultivated. Old trees are valued for their durable carpentry wood. The bark is grey and flakes off in long plates. Young branches carry felt-like hair, The leaf is a compound of 5 to 9 lanceolate leaf segments. When rubbed they give off a strong smell. The leaf edge is serrated and fimbriate. The leaf axis is pretty well bald and sometimes remains hanging on the tree for a long time in the winter. The leaves are somewhat brown tinted when they first appear but are light green in the summer and flaming yellow in the autumn. The round, sweet, edible nuts are contained in a smooth husk that opens practically completely. Deep root system.