



## Cornus controversa



<b>Výška</b>	6 – 8 (15) m
<b>Šírka</b>	6-8m
<b>Koruna</b>	round, branching in layers, half-open crown, capricious growing
<b>Kôra a vetvy</b>	green brown with shallow veins
<b>Listy</b>	wide ovate to wide oval, green, 8 – 15 cm
<b>Jesenné sfarbenie</b>	Červená, Fialová
<b>Kvety</b>	creamy-white, in umbel clusters, May/June
<b>Plody</b>	small dark-blue berry
<b>Tríne</b>	Žiadny
<b>Jedovatosť</b>	Zvyčajne nie je jedovaté pre ľudí a väčšie zvieratá
<b>Typ pôdy</b>	all, if well-drained, no dry sandy soil
<b>Spevnené plochy</b>	Netoleruje spevnené plochy
<b>Zóna odolnosti</b>	6b (-20,5 do -17,8 °C)
<b>Odolnosť voči vetru</b>	good
<b>Ostatná odolnosť</b>	Odolné mrazu (WH 1–6), Odolné vetru
<b>Stromy pre zvieratá a hmyz</b>	Odolné mrazu (WH 1–6), Odolné vetru, Plody ako potrava pre vtáctvo
<b>Použitie</b>	stromy do nádob, malé záhrady, dvory
<b>Tvar</b>	Vysokokmeň, Viackmeň
<b>Pôvod</b>	Japan, Korea, China

Small tree, its hallmark being the horizontal branches, growing in layers. In its natural surroundings it can attain a height of 15 m. The crown can be as wide as it is high. The twigs change from red brown to grey. The alternate leaf is irregular in shape and size and has a shiny dark green colour. The underside is strikingly blue green. In autumn it turns into purple red. The umbel clusters can become 15 cm across. They are short-stalked and are arranged along the horizontal branches. The many, dark-blue berries appear in autumn. In the Netherlands the fruit crop is infrequent and certainly not guaranteed. Shallow rooting with a fibrous root system. Due to an early flow of sap, pruning should take place in summer or in autumn.