



Quercus coccinea 'Splendens'



Výška	8 m
Šírka	12-15m
Koruna	flat-topped spherical, half-open crown
Kôra a vetvy	twigs brown, with thorns, bark grey-brown, slightly grooved
Listy	broad oval, glossy dark green, 5 - 10 cm, autumn colour yellow, orange and red
Jesenné sfarbenie	Červená
Kvety	white, in felt-like panicles, May-June
Plody	round, bright red, Ø about 1 cm
Tríne	Žiadny
Jedovatosť	Zvyčajne nie je jedovaté pre ľudí a väčšie zvieratá
Typ pôdy	cannot tolerate wet soil
Pôdna vlhkosť	Vhodné do vlhkých pôd
Spevnené plochy	Netoleruje spevnené plochy
Zóna odolnosti	5b (-26,0 do -23,4 °C)
Odolnosť voči vetru	good
Ostatná odolnosť	Odolné mrazu (WH 1–6), Odolné vetru, Odolné posypovej soli
Použitie	bulváre a široké ulice, priemyselné oblasti
Tvar	Vysokokmeň, Viackmeň
Pôvod	Knap Hill Nursery, Woking, England, circa 1900

Forms a broad crown with branches growing horizontally. The bark is dark grey to almost black and is moderately rough. *Q. coccinea* bears a strong resemblance to *Q. palustris*. But *Q. coccinea*'s young twigs are yellowish brown while those of *Q. palustris* are olive-green. The leaves are elongated to oval (sometimes obovate) and pinnated with 7 - 9 projecting serrated lobes. They are less deeply incised than those of *Q. palustris* and are 9 -15 cm long and 6 - 11 (15) cm wide. In the autumn the leaves turn a deep scarlet red. This *Quercus coccinea* selection is distinguished from the species by its even more intense autumnal colour. Once the leaves have turned brown they remain on the tree until halfway through winter. The acorns are single and one-third enclosed by the cupule.