



## Quercus robur 'Timuki'



Výška	15 - 20 m
Šírka	8-12m
Koruna	round, half-open crown
Kôra a vetvy	twigs dark brown, bark dark grey, deeply grooved
Listy	obovate, lobed, dark greenish purple, 7 - 14 cm
Kvety	green-yellow ? catkins, 2 - 4 cm, May
Plody	ovate to oval, 1.5 - 2.5 cm, in groups of 2 - 6, cupule with triangular contiguous scales
Tríne	Žiadny
Jedovatosť	Zvyčajne nie je jedovaté pre ľudí a väčšie zvieratá
Typ pôdy	fertile and moist
Pôdna vlhkosť	Vhodné do vlhkých pôd
Spevnené plochy	Netoleruje spevnené plochy
Zóna odolnosti	5a (-28,8 do -26,1 °C)
Odolnosť voči vetru	good
Ostatná odolnosť	Odolné mrazu (WH 1–6), Odolné vetru, Odolné posypovej soli
Použitie	priemyselné oblasti
Tvar	Vysokokmeň
Pôvod	Raimond Cinovskis, Riga, Latvia, 1946

Red-leaved cultivar of *Q. robur* with a broad ovoid crown, eventually becoming round. Grows significantly better than the old purple-leaved cultivar, 'Purpurascens'. The bark, blossom and fruit are the same as the species. The obovate leaves are, in terms of shape, also like the species although they are larger than on 'Purpurascens'. On emerging the leaf is green but it gradually turns darker, and at the end of summer it turns through red to a glossy dark greenish purple. The tree was found by M. Timuki in 1939 but was only named and distributed in 1946. The good growth and lovely foliage colours are the most important features of this purple-leaved oak. In addition, 'Timuki' is significantly less susceptible to mildew.