



Robinia ×ambigua 'Bellarosea'



Výška	6 - 8 m
Šírka	4-6m
Koruna	round, open crown, half-open crown
Kôra a vetvy	grey, deeply grooved, young shoots slightly sticky, thorny
Listy	imparipinnate, with 13 - 21 leaf segments, elliptical, green, 1 - 4 cm, feathered leaf
Kvety	in loose clusters, pink, June/August
Plody	flat skin, brown, elongated and smooth, 5 - 9 cm long
Tríne	Áno
Jedovatosť	Má jedovaté časti
Typ pôdy	places little demands on soil, provided there is good drainage
Pôdna vlhkosť	Vhodné do suchých pôd
Spevnené plochy	Toleruje spevnenie
Zóna odolnosti	3 (-40,0 do -24,5 °C)
Odolnosť voči vetru	bad
Ostatná odolnosť	Odolné mrazu (WH 1–6), Odolné posypovej soli
Stromy pre zvieratá a hmyz	Odolné mrazu (WH 1–6), Odolné posypovej soli, Cenné pre včely (medonosné)
Použitie	priemyselné oblasti
Tvar	Vysokokmeň
Pôvod	Leiden, Netherlands, approx. 1860

Robinia x ambigua is a hybrid between R. pseudoacacia and R. viscosa. It has many similarities with R. pseudoacacia but its leaves are slightly smaller. The sticky, young twigs are characteristic of R. viscosa. 'Bellarosea' is a small tree with an elegant shape. The crown is rounded and open. Unlike R. x ambigua, 'Bellarosea' has branches bearing strong thorns. The imparipinnate leaves are about 24 cm in size and consist of 13 - 21 elliptical green leaf segments. Pendulous, loose clusters of large pink flowers appear in June. The flowering period lasts into August. The branches are breakable; inclement weather conditions can cause branches to break particularly when the tree is in full bloom. Consequently, a site sheltered from the wind is required. Robinia xambigua is not very sensitive to road salt and tolerates paving well.