



Robinia viscosa



Áno

Výška	6 - 10 (12) m
Šírka	6-10m
Koruna	broad ovoid, half-open crown
Kôra a vetvy	grey, rough grooves, brownish red twigs, very sticky
Listy	imparipinnate, with 11 - 21 leaf segments, green, 2 - 5 cm, feathered leaf
Kvety	tight clusters of light pink flowers, June/July
Plody	flat pod, dark brown, very hairy, sticky, 4 - 8 cm
Tríne	Áno
Jedovatosť	Má jedovaté časti
Typ pôdy	places little demands on soil, provided there is good drainage
Pôdna vlhkosť	Vhodné do suchých pôd
Spevnené plochy	Toleruje spevnenie
Zóna odolnosti	5a (-28,8 do -26,1 °C)
Odolnosť voči vetru	sensitive
Ostatná odolnosť	Odolné mrazu (WH 1–6), Odolné posypovej soli
Stromy pre zvieratá a hmyz	Odolné mrazu (WH 1–6), Odolné posypovej soli, Cenné pre včely (medonosné)
Použitie	priemyselné oblasti
Tvar	Vysokokmeň, Viackmeň
Pôvod	eastern and south-eastern USA

Grows to heights of 12 m in its natural habitat, usually just to 8 m when cultivated. The tree has a rounded to broad ovoid crown. The many sticky parts are the most remarkable. The brownish red twigs have no thorns but are covered thickly with sticky glandular hairs. The imparipinnate leaves consist of 11 - 21 leaf segments. They are ovoid with a pointed top and are 2 - 5 cm long. They are dark green and smooth on top, and on the bottom they have grey hairs and are slightly sticky. The dark red leaf stems are also sticky. The remountant flowering starts in June and can continue into August. Light pink flowers appear in long clusters up to approx. 10 cm long. The flowers have no scent. Flat pods follow in the autumn. These are 4 - 8 cm long and covered thickly with sticky glandular hairs.