



## Tilia cordata 'Rancho'



Výška	10 - 15 m
Šírka	6-9m
Koruna	narrow conical, later narrow ovoid, half-open crown
Kôra a vetvy	dark brown, grooved trunk, young twigs green, reddish-brown later
Listy	more or less round, heart-shaped base, dark green, 5 to 12 cm
Jesenné sfarbenie	Žltá
Kvety	in umbels, 5 - 10 together, yellow to cream-coloured, pungent, June/July, fragrant flowers
Plody	round-ovoid, Ø 0.5 – 0.8 cm, first felt-like, later bare
Trne	Žiadny
Jedovatosť	Zvyčajne nie je jedovaté pre ľudí a väčšie zvieratá
Typ pôdy	any, tolerates dry conditions
Pôdna vlhkosť	Vhodné do suchých pôd
Spevnené plochy	Toleruje spevnenie
Zóna odolnosti	4 (-34,4 do -28,9 °C)
Odolnosť voči vetru	very good, also tolerates sea wind
Ostatná odolnosť	Odolné mrazu (WH 1–6), Odolné vetru
Stromy pre zvieratá a hmyz	Odolné mrazu (WH 1–6), Odolné vetru, Cenné pre motýle
Použitie	bulváre a široké ulice, úzke ulice, pobrežné zóny
Tvar	Vysokokmeň
Pôvod	E. H. Scanlon, Olmsted Falls, Ohio, USA, 1961

American cultivar that greatly resembles 'Böhlje'. Develops a tight, narrow, pyramidal crown that later becomes narrow ovoid. 'Rancho' has a regular, upwards-slanting branching pattern. The lower branches are at a 45° angle to the straight trunk; the upper branches are at a 30° angle. The tree ultimately becomes approx. 6 m wide. The dark green leaves are somewhat larger and wider than those of the species, with a heart-shaped base, tapering to an abrupt point at the top. The leaves turn a remarkable yellow colour in the autumn. Pungent flowers appear in early summer, growing in short umbels. The light green bract is somewhat shorter than the umbel. Small fruit follows in late summer. T. cordata and its cultivars are fairly unsusceptible to aphids. Its narrow manner of growth makes 'Rancho' excellently suitable for streets and avenues.