



Acer ×zoeschense 'Annae'



Výška	8 - 10 m
Šírka	6-8m
Koruna	round, spherical, half-open crown
Kôra a vetvy	bark grey-brown, young twigs red
Listy	3/5-lobed, dark green, 5 - 8 cm
Jesenné sfarbenie	Oranžová, Červená
Kvety	yellow-green corymbs, May
Plody	hairy, upright wings, 3 - 5 cm
Tŕne	Žiadny
Jedovatosť	Zvyčajne nie je jedovaté pre ľudí a väčšie zvieratá
Typ pôdy	any soils
Spevnené plochy	Toleruje čiastočné spevnenie
Zóna odolnosti	6a (-23,3 do -20,6 °C)
Odolnosť voči vetru	good
Ostatná odolnosť	Odolné mrazu (WH 1–6), Odolné vetru
Stromy pre zvieratá a hmyz	Odolné mrazu (WH 1–6), Odolné vetru, Cenné pre motýle
Použitie	stromy do nádob, strešné záhrady, priemyselné oblasti
Tvar	Vysokokmeř, Viackmeř
Pôvod	Schwerin, Germany, 1908

This medium-sized tree is a hybrid between *A. campestre* and *A. cappadocicum* subsp. *lobelii*. *A. zoeschense 'Annae'* has an open, almost spherical crown. The tree has a regular, layered habit and can reach a height of circa 10 m. The young and slender twigs that receive sufficient sunlight, turn a striking red-brown. Mature branches have a grey-brown colour. *A. zoeschense 'Annae'* does not form corky fissures on its twigs. The leaf, slightly resembling *A. campestre*, unfurls dark purple, fading gradually into dark green. The leaf margin is slightly more curvaceous than of the *A. campestre* and is a little larger. Important ornamental feature is the autumn colouring of the glossy, dark green leaf into many red to orange hues. In a sunny site it often stands out better. The fibrous root growth is shallow. Good resistance to air pollution and insensitive to diseases and pests.