



Acer triflorum



Výška	10 m
Šírka	6-8m
Koruna	round, half-open crown
Kôra a vetvy	young twigs yellowish brown, older bark brownish grey, slightly scaling
Listy	trifoliate, green, 3 - 7 cm
Jesenné sfarbenie	Oranžová, Červená, Fialová
Kvety	yellow, in smalls corymbs of three, April/May
Plody	winged nutlets, 4 - 5 cm long, hairy, 4 - 5 cm
Térne	Žiadny
Jedovatosť	Zvyčajne nie je jedovaté pre ľudí a väčšie zvieratá
Typ pôdy	neutral to slightly acidic
Spevnené plochy	Netoleruje spevnené plochy
Zóna odolnosti	6 (-23,3 do -17,8 °C)
Odolnosť voči vetru	good
Ostatná odolnosť	Odolné mrazu (WH 1–6), Odolné vetru
Použitie	stromy do nádob, strešné záhrady, malé záhrady, dvory
Tvar	Vysokokmeň, Viackmeň
Pôvod	Northern China, Korea

Bush to large tree, rarely medium-size. Somewhat resembles *A. griseum* at first sight, but differs from this in (among other things) its brown-grey, only slightly scaling bark. With *A. griseum*, the bark is reddish brown and profusely scaling. The leaves of *A. triflorum* are hairy and rather more blue-green on the underside. In summer they are green, but in autumn they turn a splendid orange to purplish red. The species name is derived from the fact that each flower corymb comprises only three flowers. *A. triflorum* should mainly be used as a solitary tree in a garden or park. It prefers a slightly acidic soil.