



Alnus glutinosa 'Laciniata'



Výška	8 - 12 m
Šírka	6-8m
Koruna	wide conical, half-open crown
Kôra a vetvy	young twigs purple grey, later grey and fissured
Listy	very deeply cut, dark green, 3 - 8 cm
Kvety	catkins, ? brown yellow, ? oval, March/April
Plody	ovate strobiles
Trne	Žiadny
Jedovatosť	Zvyčajne nie je jedovaté pre ľudí a väčšie zvieratá
Typ pôdy	moist to wet, not too poor
Pôdna vlhkosť	Vhodné do vlhkých pôd, Odolné voči krátkodobému zaplaveniu, Odolné voči dlhodobému zaplaveniu
Spevnené plochy	Toleruje spevnenie
Zóna odolnosti	3 (-40,0 do -24,5 °C)
Odolnosť voči vetru	good
Ostatná odolnosť	Odolné mrazu (WH 1–6), Odolné vetru, Odolné posypovej soli
Použitie	pobrežné zóny, priemyselné oblasti
Tvar	Vysokokmeň, Viackmeň
Pôvod	St. Germain, France, 1819

Sometimes multiple-stemmed, average-sized tree. The alternate leaves remain much smaller than of the species. The leaf is deeply lobed, slightly toothed and the lobes are lanceolate and less deep than of the 'Imperialis'. In autumn it remains long on the tree. Both the stalked buds and the young twigs are sticky to the touch (glutinosa = sticky). Female flowers are inconspicuous, male flowers in circa 7 cm long brown yellow catkins. The strobiles appear from September onwards and remain long on the tree. Besides a decorative leaf the tree also has a nice habit with slightly weeping twigs. This makes it less suitable for narrow spaces. The tree roots strongly with a branching main root. 'Laciniata' is tolerant to hard surfaces and withstands hard pruning well.