



## Robinia viscosa



Áno

<b>Výška</b>	6 - 10 (12) m
<b>Šírka</b>	6-10m
<b>Koruna</b>	broad ovoid, half-open crown
<b>Kôra a vetvy</b>	grey, rough grooves, brownish red twigs, very sticky
<b>Listy</b>	imparipinnate, with 11 - 21 leaf segments, green, 2 - 5 cm, feathered leaf
<b>Kvety</b>	tight clusters of light pink flowers, June/July
<b>Plody</b>	flat pod, dark brown, very hairy, sticky, 4 - 8 cm
<b>Tríne</b>	Áno
<b>Jedovatosť</b>	Má jedovaté časti
<b>Typ pôdy</b>	places little demands on soil, provided there is good drainage
<b>Pôdna vlhkosť</b>	Vhodné do suchých pôd
<b>Spevnené plochy</b>	Toleruje spevnenie
<b>Zóna odolnosti</b>	5a (-28,8 do -26,1 °C)
<b>Odolnosť voči vetru</b>	sensitive
<b>Ostatná odolnosť</b>	Odolné mrazu (WH 1–6), Odolné posypovej soli
<b>Stromy pre zvieratá a hmyz</b>	Odolné mrazu (WH 1–6), Odolné posypovej soli, Cenné pre včely (medonosné)
<b>Použitie</b>	priemyselné oblasti
<b>Tvar</b>	Vysokokmeň, Viackmeň
<b>Pôvod</b>	eastern and south-eastern USA

Grows to heights of 12 m in its natural habitat, usually just to 8 m when cultivated. The tree has a rounded to broad ovoid crown. The many sticky parts are the most remarkable. The brownish red twigs have no thorns but are covered thickly with sticky glandular hairs. The imparipinnate leaves consist of 11 - 21 leaf segments. They are ovoid with a pointed top and are 2 - 5 cm long. They are dark green and smooth on top, and on the bottom they have grey hairs and are slightly sticky. The dark red leaf stems are also sticky. The remontant flowering starts in June and can continue into August. Light pink flowers appear in long clusters up to approx. 10 cm long. The flowers have no scent. Flat pods follow in the autumn. These are 4 - 8 cm long and covered thickly with sticky glandular hairs.