

https://www.vdberk.sk/stromy/zanthoxylum-piperitum/



Zanthoxylum piperitum



| Výška | 3-5 m |
|----------------------------|---|
| Šírka | 3-5m |
| Koruna | round, half-open crown, capricious growing |
| Kôra a vetvy | twigs brown and short-haired, bark grey |
| Listy | imparipinnate with 11 - 23 leaf segments, oval, glossy dark green, 1.5 - 3 cm, feathered leaf |
| Jesenné sfarbenie | Žltá |
| Kvety | in axillary clusters, green, June |
| Plody | fleshy, round capsule, reddish, aromatic |
| Tŕne | Áno |
| Jedovatosť | Zvyčajne nie je jedovaté pre ľudí a väčšie zvieratá |
| Typ pôdy | few requirements |
| Spevnené plochy | Netoleruje spevnené plochy |
| Zóna odolnosti | 6 (-23,3 do -17,8 °C) |
| Odolnosť voči vetru | fairly to good |
| Ostatná odolnosť | Odolné mrazu (WH 1–6), Odolné vetru |
| Stromy pre zvieratá a hmyz | Odolné mrazu (WH 1–6), Odolné vetru, Cenné pre motýle |
| Použitie | stromy do nádob, strešné záhrady |
| Tvar | Vysokokmeň |
| Pôvod | Northern China, Korea, Japan |

Grows from a shrub shape to small, sometimes multi-stemmed tree with a round crown. The year-old twigs are short, almost wax-haired and later bare. They are brown and there are two flat thorns under each bud. The old bark is grey and sometimes the old thorns are visible as small bumps. The graceful, imparipinnate leaves are 7 - 15 cm long and consist of 11 to 23 oval leaf segments. The margins are finely serrated and the tip of the terminal leaflet is dissected to retuse. The leaf stem is hairy and has a few small prickles on the underside. The leaf turns yellow in autumn. Compact clusters of green flowers open in June. They are followed by small reddish spherical fleshy capsules. These contain black seeds. In Japan the seeds are ground and used as pepper. A park tree that has few soil requirements.